Vivostat[®] Co-Delivery

By using the Vivostat[®] Co-Delivery system it is possible to co-deliver cells, drugs etc. with Vivostat[®] Fibrin Sealant or Vivostat PRF[®] (Platelet Rich Fibrin)





Vivostat[®] Co-Delivery

Vivostat has developed the revolutionary Co-Delivery system that makes it possible to co-deliver a desired substance¹ (drugs, cells etc.) with Vivostat[®] Fibrin Sealant or Vivostat PRF[®] (Platelet Rich Fibrin)

The opportunities with the Vivostat[®] Co-Delivery system are vast and the system allows the surgeon to apply a selected substance easily and effectively. Furthermore, it may be possible to reduce the total cost of a procedure by using the Vivostat[®] Co-Delivery system².

Options for Co-Delivery include:

bone allograft for posterolateral multilevel cervical fusion

Drugs

- Antimicrobials
- Chemotherapeutics
- Pain medications

Cells

- Stem cells
- Skin cells

Co-delivering drugs, cells etc. with the Vivostat[®] Fibrin Sealant or Vivostat PRF[®] solutions offers the surgeon and the patient a number of benefits:

- Topical application
- Targeting affected/desired area
- Possible higher local dose
- Possible lower systemic impact
- Improved compliance

Moreover, no thrombin is added to Vivostat[®] Fibrin Sealant and Vivostat PRF[®] (unlike most other sealants and PRP products). This is beneficial to the Co-Delivery system as thrombin activation has been shown to have a negative effect on cell survival³.

The fibrin membrane found in both Vivostat[®] Fibrin Sealant and Vivostat PRF[®] has, furthermore, been shown to postpone the degradation process of the substance. This means that the fibrin membrane ensures a slow and sustained release of the substance offering a prolonged effect⁴.


How does it work

Up to 5 ml of substance may be co-delivered together with the Fibrin or PRF® solution. The substance is applied using the Vivostat® Spraypen⁵ which enables the surgeon to apply the substance accurately and intermittently throughout the entire procedure. The substance and the Fibrin or PRF® solution is mixed once it leaves the tip of the Spraypen and polymerizes immediately upon application - this way the substances stay where they are intended to act.

Vadalà et al. - Journal of Tissue Engineering and Regenerative Medicine 2008; 2: 515–520.

tration in an immune-competent rat model with malignant pleuromesothelioma Lardinois et al. - J Thorac Cardiovasc Surg 2006;131:697-703

5) It can also be applied endoscopically



The Vivostat[®] Co-Delivery tube (Cross-section)



Co-Delivery Leaflet ver. 2

The Vivostat[®] Spraypen is used together with the Vivostat[®] foot switch, and the APL 404 to perform the Co-Delivery application.

ponsibility of the surgeon.
 3) High thrombin concentrations in fibrin sealants induce apoptosis in human karatinocytes

 etrich fibrin on corticocancellous
 3 High thrombin concentrations in fibrin sealants induce apoptosis in human karatinocytes

 etrich fibrin on corticocancellous
 3 High thrombin concentrations in fibrin sealants induce apoptosis in human karatinocytes

 etrich fibrin on corticocancellous
 3 High thrombin concentrations in fibrin sealants induce apoptosis in human karatinocytes

 etrich fibrin on corticocancellous
 4 Intrapleural topical application of cisplatin with the surgical carrier //vostat increases the local drug concenter